ISSN (print) 0093-4666 © 2013. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/125.91

Volume 125, pp. 91–96 *July–September 2013*

Two new species of *Septobasidium* (*Septobasidiaceae*) from Yunnan Province in China

Wei Li[1,2] & Lin Guo[1*]

[1]*State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*

[2]*Graduate University of Chinese Academy of Sciences, Beijing 100049, China*

* Correspondence to: *guol@im.ac.cn*

Abstract —Two new species, *Septobasidium fissuratum* on *Castanea* sp. associated with *Pseudaulacaspis* sp. and *Septobasidium heliciae* on *Helicia nilagirica* associated with *Lepidosaphes* sp., are described.

Key words —*Pucciniomycetes*, *Septobasidiales*, taxonomy

Two new species of *Septobasidium* were recently discovered from Yunnan Province. These specimens, collected by Dr. Shuanghui He, are described below.

Septobasidium fissuratum Wei Li bis & L. Guo, **sp. nov.** Figs 1–7

Fungal Name FN 570061

Differs from *Septobasidium meridionale* by its larger basidia.

Type: China, Yunnan Province, Longchuan, Longba, alt. 1000 m, on *Castanea* sp. (*Fagaceae*), associated with *Pseudaulacaspis* sp. (*Diaspididae*), 31.X.2012, S.H. He YN08 (HMAS 244419, **holotype**).

Etymology: The epithet refers to the fissures of basidiomata.

Basidiomata on branches, 10–14 cm long, 2.5–6 cm wide, pale cinnamon-brown or cinnamon-brown; margin determinate, surface smooth, with many fissures at maturity. In section 600–1100 μm thick, composed of 3–4 layers. Subiculum brown, 30–50 μm thick. Pillars brown, 100–180 μm high, 30–100 μm wide, branched outwards to form a hyphal layer 350–700 μm high, then forming 20–40 μm thick horizontal layer, from which the fungal hyphae successively producing hyphal layers at maturity. Hymenium hyaline or brownish, 60–100



Fig. 1. *Septobasidium fissuratum* (HMAS 244419, holotype). Basidia.

μm thick. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or slightly curved, 4-celled, 32–45 × 6–9 μm, hyaline or brownish. Basidiospores not seen. Haustoria consisting of irregularly coiled hyphae.

Comments: *Septobasidium fissuratum* is similar to *S. meridionale* C.X. Lu & L. Guo, but the latter has fewer fissures on the basidiome surface, an inconspicuous horizontal layer in the section, and smaller basidia, 27–36 × 7–9.5 μm (Lu & Guo 2010b).

Septobasidium heliciae Wei Li bis & L. Guo, **sp. nov.** Figs 8–14

Fungal Name FN 570062

Differs from *Septobasidium septobasidioides* by its wider pillars and smaller basidia.

Type: China, Yunnan Province, Yingjiang, Tongbiguan Nat. Res., alt. 1250 m, on *Helicia nilagirica* (*Proteaceae*), associated with *Lepidosaphes* sp. (*Diaspididae*), 29.X.2012, S.H. He YN07 (HMAS 244418, **holotype**).

ETYMOLOGY: The epithet refers to the substrate plant genus.

Basidiomata on branches, 4–13 cm long, 4–5 cm wide, white or pale cinnamon-brown; margin determinate, surface smooth. In section 750–1100 μm thick.

Figs. 2–7 *Septobasidium fissuratum* (HMAS 244419, holotype). 2. Basidiomata on branch. 3–4. Sections of basidiomata (arrow shows horizontal layer). 5–6. Basidia (arrows). 7. Haustoria.


2
1 cm


3
100 µm


4
100 µm


5
10 µm


6
10 µm


7
10 µm



Fig. 8. *Septobasidium heliciae* (HMAS 244418, holotype). Basidia.

Subiculum brown, 10–20 μm thick. Pillars brown, 120–600 μm high, 30–120 μm wide, branched outwards to form a hymenial layer. From the old hymenium successively forming new hyphal layer or pillars and hymenium. Hymenium hyaline or brownish, 80–110 μm thick, with irregularly arranged hyphae. Basidia arising directly from the hyphae without a probasidial cell, cylindrical, straight or curved, 4-celled, 30–45 × 5–6.5 μm, hyaline or brown. Basidiospores not seen. Haustoria consisting of coiled hyphae.

Comments: *Septobasidium heliciae* is similar to *S. septobasidioides* (Henn.) Höhn. & Litsch., but the latter has basidiomata with lobed wings, slender pillars (10–25 μm wide), and larger basidia (40–55 × 8.4–10 μm; Couch 1938).

Including the two new species reported in this paper, 51 *Septobasidium* species have now been reported in China (Sawada 1933, Couch 1938, Teng 1963, Tai 1979, Kirschner & Chen 2007, Lu & Guo 2009a,b,c, 2010a,b,c, 2011, Lu et al. 2010, Chen & Guo 2011a,b,c,d, 2012a,b,c).

Figs. 9–14. *Septobasidium heliciae* (HMAS 244418, holotype). 9. Basidiomata on branches. 10–11. Sections of basidiomata. 12–13. Basidia. 14. Haustoria.


9
1 cm

10
100 µm


11
100 µm


12
10 µm


13
10 µm


14
10 µm

Acknowledgements

The authors would like to express their deep thanks to Dr. Eric H.C. McKenzie (Auckland, New Zealand) for serving as pre-submission reviewer, to Dr. Shuanghui He (Beijing Forestry University) for serving as pre-submission reviewer and collecting specimens, to Dr. Shaun Pennycook (Auckland, New Zealand) for nomenclatural review, to Prof. Zhenyu Li (Institute of Botany, Chinese Academy of Sciences) for identifying the host plants, to Prof. Sanan Wu (Beijing Forestry University) for identifying the scale insects, and to Mrs. Xiangfei Zhu for inking in line drawings. This study was supported by the National Natural Science Foundation of China (No. 30670005).

Literature cited

Chen SZ, Guo L. 2011a. *Septobasidium sichuanense* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 115: 481–484. http://dx.doi.org/10.5248/115.481

Chen SZ, Guo L. 2011b. *Septobasidium atalantiae* sp. nov. (*Septobasidiaceae*) and *S. henningsii* new to China. Mycotaxon 117: 291–296. http://dx.doi.org/10.5248/117.291

Chen SZ, Guo L. 2011c. *Septobasidium saurauiae* sp. nov. (*Septobasidiaceae*) and *S. pseudopedicellatum* new to China. Mycotaxon 118: 283–288. http://dx.doi.org/10.5248/118.283

Chen SZ, Guo L. 2011d. *Septobasidium glycosmidis* and *S. albiziae* spp. nov. (*Septobasidiaceae*) from Hainan Province. Mycosystema 30: 861–864.

Chen SZ, Guo L. 2012a. Three new species and three new Chinese records of *Septobasidium* (*Septobasidiaceae*). Mycosystema 31: 651–655.

Chen SZ, Guo L. 2012b. Three new species of *Septobasidium* (*Septobasidiaceae*) from Hainan Province in China. Mycotaxon 120: 269–276. http://dx.doi.org/10.5248/120.269

Chen SZ, Guo L. 2012c. Three new species of *Septobasidium* (*Septobasidiaceae*) from southern and southwestern China. Mycotaxon 121: 375–383. http://dx.doi.org/10.5248/121.375

Couch JN. 1938. The genus *Septobasidium*. Univ. of North Carolina Press, Chapel Hill. 480 p.

Kirschner R, Chen CJ. 2007. New reports of two hypophyllous *Septobasidium* species from Taiwan. Fung. Sci. 22(1,2): 39–46.

Lu CX, Guo L. 2009a. *Septobasidium maesae* sp. nov. (*Septobasidiaceae*) from China. Mycotaxon 109: 103–106. http://dx.doi.org/10.5248/109.103

Lu CX, Guo L. 2009b. Two new species of *Septobasidium* (*Septobasidiaceae*) from China. Mycotaxon 109: 477–482. http://dx.doi.org/10.5248/109.477

Lu CX, Guo L. 2009c. *Septobasidium annulatum* sp. nov. (*Septobasidiaceae*) and *Septobasidium kameii* new to China. Mycotaxon 110: 239–245. http://dx.doi.org/10.5248/110.239

Lu CX, Guo L. 2010a. Three new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. Mycotaxon 112: 143–151. http://dx.doi.org/10.5248/112.143

Lu CX, Guo L. 2010b. Two new species of *Septobasidium* (*Septobasidiaceae*) and *S. pallidum* new to China. Mycotaxon 113: 87–93. http://dx.doi.org/10.5248/113.87

Lu CX, Guo L. 2010c. Two new species of *Septobasidium* (*Septobasidiaceae*) from Hainan province in China. Mycotaxon 114: 217–223. http://dx.doi.org/10.5248/114.217

Lu CX, Guo L. 2011. Two new species of *Septobasidium* (*Septobasidiaceae*) from Gaoligong Mountains in China. Mycotaxon 116: 395–400. http://dx.doi.org/10.5248/116.395

Lu CX, Guo L, Wei JG, Li JB. 2010. Two new species of *Septobasidium* (*Septobasidiaceae*) from southern China. Mycotaxon 111: 269–274. http://dx.doi.org/10.5248/111.269

Sawada K. 1933. Descriptive catalogue of the Formosan fungi. Part VI. Rep. Dept. Agric. Govt. Res. Inst. Formosa 61: 1–99.

Tai FL. 1979. Sylloge Fungorum Sinicorum. Science Press, Beijing. 1527 p.

Teng SC. 1963. Fungi of China. Science Press, Beijing. 808 p.